

# **IBM**

## **Exam C2020-011**

### **IBM SPSS Statistics Level 1 v2**

**Version: 6.0**

**[ Total Questions: 55 ]**

**Question No : 1**


What statistical test should be used to assess whether the percentage differences observed in a crosstabs table could have occurred by chance?

- A. Correlation
- B. Linear Regression
- C. T-Test
- D. Chi-square test of independence

**Answer: D**

**Question No : 2**

The text file shown in the figure below is an example of a fixed format text file.



- A. True
- B. False

**Answer: B**

**Question No : 3**

In the Variable View, if you have a series of variables that share the same category coding scheme, you can enter value labels for one variable, then copy these labels to the other variables.

- A. True
- B. False

**Answer: A**

**Question No : 4**

For a variable salary we have the statistics as shown in the figure below.

	N	Minimum	Maximum	Mean
Current Salary	474	-9999.00	135000.00	33916.7321
Valid N (listwise)	474			


The reported Mean is incorrect because a value, -9999, is included in its calculation. How can this situation be fixed?

- A. Define -9999 as a system-missing value.
- B. Remove allcases with the value -9999 for the variable salary from the data file.
- C. Define -9999 as a user-missing value.
- D. Define -9999 as both a system- and user-missing value.

**Answer: C**

**Question No : 5**

Which statement is true about this box plot?



- A. The mean is 14.
- B. The standard deviation is 14 (20 minus 6).
- C. The standard deviation is 4 (16 minus 12).
- D. of the outliers are on the lower end of the distribution.

Answer: D

**Question No : 6**

Which statement is the correct interpretation of this crosstab table?

BELIEF IN LIFE AFTER DEATH \* GENDER OF RESPONDENT Crosstabulation

			GENDER OF RESPONDENT		Total
			Female	Male	
BELIEF IN LIFE AFTER DEATH	YES	Count	541	417	958
		% within BELIEF IN LIFE AFTER DEATH	56.5%	43.5%	100.0%
		% within GENDER OF RESPONDENT	86.0%	76.9%	81.8%
	NO	Count	88	125	213
		% within BELIEF IN LIFE AFTER DEATH	41.3%	58.7%	100.0%
		% within GENDER OF RESPONDENT	14.0%	23.1%	18.2%
Total	Count	629	542	1171	
	% within BELIEF IN LIFE AFTER DEATH	53.7%	46.3%	100.0%	
	% within GENDER OF RESPONDENT	100.0%	100.0%	100.0%	

- A. 56.5% of females believe in life after death.
- B. 86.0% of females believe in life after death.
- C. 27.5% of females believe in life after death.
- D. 53.7% of females believe in life after death.

**Answer: B**

### Question No : 7

Consider the data file below and answer the following: The calculation of the mean of the variables X, Y, Z is contained in the variable Av\_XYZ. What method was used to calculate the mean?

	X	Y	Z	Av_XYZ
1	4.00	1.00	4.00	3.00
2	.	1.00	.	.
3	4.00	2.00	3.00	3.00
4	5.00	.	6.00	5.50
5	6.00	12.00	7.00	8.33

- A. The Compute Variable dialog and the expression  $(X+Y+Z) / 3$
- B. The Compute Variable dialog and the expression  $X+Y+Z/3$
- C. The Compute Variable dialog and the expression  $MEAN(X, Y, Z)$
- D. The Compute Variable dialog and the expression  $MEAN.2(X, Y, Z)$

Answer: D

### Question No : 8

If you request an aggregated dataset or data file in the Aggregate procedure, the number of cases in the new aggregated file is equal to what?

- A. Number of cases in the original data file
- B. Number of aggregated summary variables
- C. Number of categories of the variables specified in the Break Variables list
- D. Number of cases that you specified in the Aggregate Data dialog box

Answer: C

### Question No : 9